ISyE 3770 Assignment 7: Hypothesis Testing and Linear Regression

Due date: 11:59 PM, Friday, April 26, 2024.

Question 1 (Find type-I/type-II errors and *p*-value). A manufacturer is interested in the output voltage of a power supply used in a PC. Output voltage is assumed to be normally distributed, with standard deviation 0.25 volt, and the manufacturer wishes to test $H_0: \mu = 5$ volts against $H_1: \mu \neq 5$ volts, using n = 8 units.

- 1) The acceptance region is $4.85 \le \bar{x} \le 5.15$. Find the value of α (Type I error).
- 2) Find the power of the test for detecting a true mean output voltage of 5.1 volts.
- 3) Find the p-value of the observed statistic is (i) $\bar{x} = 5.2$ and (ii) $\bar{x} = 4.7$, respectively.

Question 2 (Hypothesis testing for sample mean difference). A computer scientist is investigating the usefulness of two different design languages in improving programming tasks. Twelve expert programmers, familiar with both languages, are asked to code a standard function in both languages, and the time (in minutes) is recorded. It is assumed that the standard deviation of the first design language is 13 and that of the second is 21. Both design languages follow normal distribution. The data is presented as follows:

Design Language 117, 16, 21, 14, 18, 24, 16, 14, 21, 23, 13, 18Design Language 218, 14, 19, 11, 23, 21, 10, 13, 19, 24, 15, 20

- 1) Find a 95% confidence interval on the difference in mean coding times. Is there any indication that one design language is preferable?
- 2) If hypothesis test is performed to answer the above question: whether one design language is preferable or not. Write down the null and alternative hypothesis. Report the p-value and draw a conclusion (either accept or reject H_0) under significance level 0.05.

Question 3 (Programming Problem). Cloud seeding has been studied for many decades as a weather modification procedure (for an interesting study of this subject, see the article in Technometrics, "A Bayesian Analysis of a Multiplicative Treatment Effect in Weather Modification," Vol. 17, pp. 161 - 166). The rainfall in acre-feet from 23 clouds that were selected at random and seeded with silver nitrate is given below:

49.44, 11.33, 12.33, 30.30, 30.50, 35.40, 36.40, 34.76, 33.21, 39.66, 37.53, 31.33

36.54, 38.54, 27.32, 26.43, 25.43, 26.65, 43.33, 42.33, 44.32, 41.32, 54.33

(Hint: you can load the data using the rain.txt file in the attachment).

It is assumed that the rain fall follows normal distribution with known standard deviation $\sigma = 10$. Now, answer the following questions:

- 1) Suppose we want to test the claim that the mean rain fall is above 30. What is the null and alternative hypotheses that we should use to test this claim?
- 2) Perform a Z-test (see page 26 in slides Week11.pdf) to test the claim in question 3. Use a confidence level of 95%. What are your conclusions?

You may use:

```
z.test(data, alternative = xxx, mu=xxx, sigma.x=xxx, conf.level=xxx)
```

Where,

data should be Rain,

```
alternative is either alternative=c("greater"), or alternative=c("less")
or alternative=c("two.sided") depending on your alternative hypothesis,
```

mu should be 30,

sigma.x should be 10,

conf.level should be 0.95

3) What was the p-value for the hypothesis test in part 2)?

Question 4 (Computation in simple linear regression). Regression analysis were used to analyze the data from a study investigating the relationship between roadway surface temperature (°F) (x) and pavement deflection (y). Summary quantities were n = 20, $\sum y_i = 12.75$, $\sum y_i^2 = 8.86$, $\sum x_i = 1478$, $\sum x_i^2 = 143,215.8$, $\sum x_iy_i = 1083.67$.

- 1) Calculate the least squares estimate of the slope and intercept. Graph the regression line. Estimate σ^2 .
- 2) Use the equation of the fitted line to predict what pavement deflction would be observed when the surface temperature is 85F.
- 3) What is the mean pavement deflection when the surface temperature is 90F.
- 4) What change in mean pavement deflection would be expected for a 1F change in surface temperature.

Question 5 (Simple Linear Regression). We have data on the quarterback ratings for the 2008 National Football League season (Source: The Sports Network). It is suspected that the rating y is related to the average number of yards gained per pass attempt (x).

Getting the Data: Data in this question is contained in file data113.csv. Once you have saved the data file in the working directory, read the data in R using the command

data = read.csv("data113.csv",header=TRUE)

We can investigate the association of Rating to Yds using linear regression. Define first

y = data\$Rating

x = data\$Yds

The function to fit a linear regression model in R is lm. We perform a linear regression with R as follows

```
model = lm(y^x)
```

summary(model)

You can use the fitted results to help you answer above questions. Make sure to include the R output in your homework and state clearly how do you use the R output to solve the following questions.

- 1) Calculate the least squares estimates of the slope and intercept. What is the estimate of σ^2 ? Graph the regression model.
- 2) Find an estimate of the mean rating if a quarterback averages 7.5 yards per attempt.
- 3) What change in the mean rating is associated with a decrease of one yard per attempt?
- 4) To increase the mean rating by 10 points, how much increase in the average yards per attempt must be generated?
- 5) Find 95% confidence intervals on the slope and intercept.(Hint: you can get these CIs using R: confint (model))
- 6) Find a 95% prediction interval on the rating when the average yards per attempt is 8.0.
 (Hint: you can get these CIs using R: predict (model, data.frame(x=8), interval = c("prediction"))).

Question 6 (Multiple Linear Regression). The electric power consumed each month by a chemical plant to thought to be related to the average ambient temperature x_1 , the number of days in the month x_2 , the average product purity x_3 , and the tons of product produced x_4 . The past year's historical data are available and are presented in the following table.

у	x_1	<i>x</i> ₂	x_3	x_4
240	25	24	91	100
236	31	21	90	95
270	45	24	88	110
274	60	25	87	88
301	65	25	91	94
316	72	26	94	99
300	80	25	87	97
296	84	25	86	96
267	75	24	88	110
276	60	25	91	105
288	50	25	90	100
261	38	23	89	98

Use R to find the multiple linear regression model. For example, you may create the variable x_1 in R using the following command

x1 = c(25, 31, 45, 60, 65, 72, 80, 84, 75, 60, 50, 38)

You can fit a multiple linear regression model similar to simple linear regression using the following command

```
model = lm(y^x1+x2+x3+x4)
```

```
summary(model)
```

Based on the results or *R*, answer the following questions:

- 1) Fit a multiple linear regression model to these data.
- 2) Estimate σ^2 .
- 3) Compute the standard errors of the regression coefficients. Are all of the model parameters estimated with the same precision? Why or why not?
- 4) Predict the power consumption for a month in which $x_1 = 75$ F, $x_2 = 24$ days, $x_3 = 90\%$, and $x_4 = 98$ tons.